



Supplement of

HTAP3 Fires: towards a multi-model, multi-pollutant study of fire impacts

Cynthia H. Whaley et al.

Correspondence to: Cynthia H. Whaley (cynthia.whaley@ec.gc.ca)

The copyright of individual parts of the supplement might differ from the article licence.

Supplement of “HTAP3 Fires: Towards a multi-model, multi-pollutant study of fire impacts” by Whaley et al

Contents

Text S1: Regional discussions	1
Text S2: Health impacts from acute exposure to fire emissions	3
Text S3: Fire management policies	4
Text S4: Inverse modelling	6
Table S1: Observations for model evaluation	8
Table S2: Historical meteorological reanalysis options	10
Table S3: Health risk assessment tools	10
Table S4: Comparative analysis of health risk tools	11
References	12

S1. Regional discussions

Additional information about biomass burning in several global regions are included below to supplement that in the main paper.

Sub-Saharan Africa

Fire-induced BC concentration and direct climate forcing vary significantly spatially and temporarily, with the highest emissions globally occurring in sub-Saharan Africa (van der Werf et al., 2010). Flaming grassland fires produce higher BC emission fractions than is typical for the northern hemisphere (Dobracki et al. 2024). The impact of increased warming is already detectable on the longrange transport of BB emissions (Tatro and Zuidema, 2024).

Amazon Basin

Occurrences of wildfires in the Amazon Basin respond strongly to climate change and play an increasingly important role in changing the global climate.

Arctic

During extreme fire years like 2020 in the Arctic and boreal driven by both lightning and human-caused fires, BC emissions from wildfires as estimated by GFAS are several times larger than anthropogenic sources estimated from the GAINS model (McCarty et al., 2021). The BC levels and climate forcing sourced from wildfire biomass burning across the Northern boreal forest have increased in the past decades, particularly in recent several years (Liu et al., 2023).

The biogeochemistry in the Arctic Ocean has been shown to be influenced by Siberian fires emissions (Section 3.2.3), as well as a large fraction of the nitrogen was likely to be coming from Boreal peat burning; linking the impacts of a warming climate on fire activity to widescale ecosystem change across the Arctic region.

Australia

The 2019-2020 Australian wildfires were the most expensive to-date for that country with total costs of US\$4.5 billion (Masters, 2021).

Equatorial Asia

An estimated 100,000 premature deaths across Equatorial Asia occurred due to the 2015 severe wildland fire season in Indonesia (Kopplitz et al., 2016; Crippa et al. 2016).

Europe

In Greece, 800 families had lost their forest-based livelihoods in the wildfires on the island of Evia in 2021 (Psaropoulos, 2021). There have also been increased civilian fatalities since the late 1970s in per-urban or wildland-urban interface (WUI) areas of the Mediterranean (Molina-Terres et al. 2019). In another study of 2009, 323 896 ha were burned in France, Portugal, Spain, Greece and Italy (European commission, 2010). While Mediterranean fires are small compared to fires in other areas of the world, they often cause significant damage to homes and businesses and threaten human beings and animals, because they occur so close to urban areas and are fueled by hot temperatures and high wind speeds.

India

Forests in India are of diverse types but much focus remains on fires in Himalayas or shifting cultivation led fires in north-east India. However, central India forest fires often go neglected. In the study conducted by Jain et al. (2021), from 2001 to 2020, about 70% of forest fires over central India domain occurred during 2 months March (1,857.5 counts/month) and April (922.8 counts/month). However, owing to warmer conditions in the Indian subcontinent from 2006 to 2020 (as compared to 2001–2005), a doubling and tripling of forest fire activity is noted in forest fire (FMAMJ) and non-fire (JASONDJ) seasons, respectively. The study further highlights the role of multiple simultaneous climate extremes for example, El Niño, heat waves, weak ISM, and droughts in causing anomalously high fire activity periods over central India. Statistical associations among monthly forest fire counts and various meteorological and environmental variables over a smaller but a high forest fire activity region are highlighted for FMAMJ. High forest fire counts decrease the soil moisture content, evapotranspiration, and normalized difference vegetation index (NDVI) and are associated with an increase in BC and CO emissions. High surface air temperatures prior to ISM significantly increase soil temperature and significantly reduce NDVI, creating a drier environment. Moreover, precipitation shows a significant positive correlation with soil moisture content, evapotranspiration and NDVI. During FMAMJ, chances of precipitation are very low. Thus, high near surface air low precipitation, low soil temperature and moisture, low evapotranspiration and low NDVI during FMAMJ create conducive conditions for high forest fire activity in central India domain.

Agricultural burning smoke has been shown to cause asthma, chronic obstructive pulmonary disease (COPD), bronchitis, reduced lung capacity, emphysema, cancer, and more (Ghosh et al., 2019). For example, people residing in Punjab, agricultural state of India, are exposed to recurrent fires, and have suffered from those respiratory diseases (Alexaki et al. 2019; Saxena et al. 2021). Among these sources, agricultural crop residue burning (ACRB) stands out as a significant contributor to ambient PM, alongside biowaste burning and forest fires (Mittal et al., 2009). Stubble burning has been linked to a 3.25% increase in mortality (1.56% per month) with a 10 µg/m³ rise in particulate matter, notably leading to a 1.82% surge in cardiopulmonary disease cases (Liu et al., 2018). A mere 10-point increase in stubble burning could elevate overall mortality by 1.71% and cardiopulmonary mortality by 1.91%. The burning of crop residues in northwest India has directly impacted Delhi's Air Quality Index (AQI), contributing to 149 million tons (Mt) of carbon dioxide (CO₂), 9.0 Mt of carbon monoxide (CO), 0.25 Mt of sulphur oxide (SOX), 1.28 Mt of particulate matter (PM), and 0.07 Mt of black carbon (BC) (He et al., 2020). Heart disease affected 84.5% of the population, eye irritation impacted 76.8%, nasal irritation affected 44.8%, and throat irritation impacted 45.5%. Instances of hacking-related issues increased by 41.6%, while wheezing problems rose by 18.0% (Yadav, 2019).

Lower Mekong Region

Upper Southeast Asia has a tropical-subtropical climate and experiences fires recurring during the dry season (November-April). Its first half is influenced by the northeast monsoon, which induces cool dry air with no or little precipitation favorable for fuels to dry and fires to spread. In the second half, the monsoon weakens and ceases, with March-April corresponding to the summer when relatively high temperature promotes fire ignitability. Almost all fires in the region are caused by or linked to human activities, such as gathering of non-timber forest products, hunting, deforestation, land clearing, slash-and-burn agriculture, and carelessness (Tanpipat et al. 2009). According to Phoo et al. (2024), three distinct fire-prone areas exist in Upper Southeast Asia, which are western Myanmar, northern Cambodia, and the tri-state zone (including eastern Myanmar, northern Thailand, and northern Laos combined). The region that aggregately encompasses the last two fire-prone areas is here referred to as the Lower Mekong Region. The peak fire timing of northern Cambodia is January-February but shifted later to March-April for the other fire-prone areas. The tri-state zone has diverse land covers with forests being dominant (followed by croplands and then urbanized areas) and is also topographically complex with mountains-valleys, highlands, and plains. Pollutants emitted from fires and other anthropogenic sources can thus easily accumulate under stagnant air with low mixing height. This fire-prone area has been recognized as having severe smoke haze occurrences, which cause public health concern, socio-economic impacts, reduced visibility for transport and tourism, and transboundary air pollution (Bran et al. 2022; Moran et al. 2019; Thao et al. 2022). Effort has been made through Trilateral Consultation on Transboundary Haze Pollution, among leaders of Laos, Myanmar and Thailand, with collaboration towards making CLEAR Sky Strategy Joint-Plan of Implementation to prevent, mitigate and manage open fires (Unapumnuk and Yensrong, 2024), while scientific support is required, as emphasized in the ASEAN Haze Free Roadmap 2023-2030, starting from Thailand collaboration with NASA and International community in the field campaign conducted during March 2024 (https://espo.nasa.gov/asia-aq/content/ASIA-AQ_White_Paper).

United States

In California, along with the neighbouring states of Oregon and Washington (ODF, 2020), the last 5 years have seen a historic increase in wildfire activity. In 2017, the Tubbs Fire became the most catastrophic wildfire on state record, claiming 5,636 structures and 22 lives (CAL FIRE, 2020). The approximated economic impacts from lost infrastructure and homes, health costs and other damages in California, from overall economic and business disruptions was US\$148.5 billion in 2018, or roughly 1.5% of the state's GDP (Wang et al. 2021).

S2. Health impacts from acute exposure to fire emissions

Some VOCs and PAHs released during wildfires causes skin and eye irritation, drowsiness, coughing and wheezing while other like benzene may be carcinogenic (Sahu and Saxena, 2015). There were studies conducted in firemen provide an accurate model for understanding health impact of wildfire emissions. Extensive research conducted during forest firefighting in the United States and Australia identified carbon monoxide and respiratory irritants as the main wildfire pollutants of concern to firefighters (Aisbett et al., 2012). Brotherhood et al. (1990) assessed the carboxyhemoglobin saturation (COHb%) levels from alveolar CO levels in 24 firefighters working with hand tools and in 12 accompanying scientific observers, before and after firefighting (duration 37 to 187 min) on 15 experimental bushfires. Carboxyhemoglobin levels increased on average by 0.7% per hour in the firefighters and by 0.3% per hour in the observers that indicate that firefighters are generally unlikely

to experience hazardous levels of CO exposure. A risk assessment of firefighter exposure during wildfires was conducted in Australia (Reisen et al. 2011). This study monitored air toxins within the breathing zone of the firefighters and showed that 30% of firefighters were exposed to high levels of hazardous pollutants like CO, respirable particles and formaldehyde that increased the occupational exposure standard (OES) for 5% to 20% of time.

Similarly, farmers and those close to recurrent agricultural burning can also be acutely exposed to the smoke, with negative consequences. For example, due to prolonged pollutant presence from stubble burning and limited dispersion owing to atmospheric conditions, the smoke accumulates in the lower atmosphere, posing heightened risks to people's well-being (Govardhan et al., 2023). Prolonged exposure at high levels can result in permanent health issues, including the development of chronic lung diseases such as asthma, chronic obstructive pulmonary disease (COPD), bronchitis, reduced lung capacity, emphysema, cancer, and more (Ghosh et al., 2019). Many farmers exposed to stubble smoke report eye and lung irritation and significant healthcare expenses (Kumar et al., 2015).

Due to prolonged pollutant presence from stubble burning and limited dispersion owing to atmospheric conditions, the smoke accumulates in the lower atmosphere, posing heightened risks to people's well-being (Govardhan et al., 2023). Prolonged exposure at high levels can result in permanent health issues, including the development of chronic lung diseases such as asthma, chronic obstructive pulmonary disease (COPD), bronchitis, reduced lung capacity, emphysema, cancer, and more (Ghosh et al., 2019). Many farmers exposed to stubble smoke report eye and lung irritation and significant healthcare expenses (Kumar et al., 2015).

Finally, high blood pressure is a primary risk factor for non-communicable disease-related deaths across diverse economic settings (IHME, 2017). Medical and epidemiological investigations have connected air pollution from fires to elevated rates of hypertension and cardiovascular stress (Hadley et al., 2018). The adverse effects of exposure span from skin and eye irritation to severe neurological, cardiovascular, and respiratory ailments (Lelieveld et al., 2015). In specific cases, it can lead to fatal consequences, particularly among individuals with pre-existing respiratory conditions (Saggu et al., 2018).

S3. Fire Management Policies

S.3.1: Agricultural fire management

Amidst the escalating detrimental health effects of crop residue burning, the implementation of sustainable residue management practices is essential to safeguard both the environment and public health. Several studies have highlighted various environmentally sustainable crop residue management methods that not only offer economic advantages but also reduce healthcare expenses (Jiang et al., 2018). Alternative approaches proposed by researchers involve utilizing stubble for purposes such as raw material in liquor production, mushroom cultivation, or as fuel for boilers via gasification processes (Gummert et al., 2020). Moreover, employing crop residue in the construction industry for making diverse types of cement and blocks presents a feasible strategy (Bories et al., 2015). Despite extensive research confirming the positive impacts of Sustainable Crop Residue Management Practices (SCRMPs) and their feasibility at a local level, their adoption rate

remains relatively low (Smith et al., 2007). This limited adoption might stem from inadequate awareness about the advantages of these strategies, particularly concerning their environmental and health implications (Domingo et al., 2014).

S.3.2: Prescribed burns

In the field of conservation of biodiversity, controlled burning is an effective method. Controlled burns, which replicate natural fire regimes, provide different habitat structures that sustain a wide range of plant and animal species (Pausas and Keeley, 2019). Fire-adapted flora survives, maintains ecological balance, and avoids species domination, all of which contribute to biodiversity enhancement (He et al., 2019). Prescribed burning increases ecosystem resilience by proactively lowering fuel loads, resulting in fire-adapted landscapes that recover faster from disturbances (Thacker et al., 2023; Volkova et al., 2021). This technique not only guards against catastrophic fires, but it also increases adaptive capacity, hence supporting long-term ecological stability (Ascoli et al., 2023; Pais et al., 2023).

The reaction of plants to targeted fires is noteworthy, triggering systems that promote regeneration (Sagra et al., 2019). Fire-adapted plant species thrive thanks to adaptations including serotinous cones and fire-resistant bark (He et al., 2019; Pausas and Keeley, 2019). Controlled fires also eliminate invasive species and stimulate the germination of certain seeds, modifying vegetation composition in accordance with ecological goals (Ditomaso et al., 2006; Lamont, 2022; Sagra et al., 2019).

Addressing air quality concerns is a critical component of sustainable land management. Although controlled fires emit emissions, planned execution reduces their negative effects (Bowman et al., 2018; Jaffe et al., 2020; Neidermeier et al., 2023). Timing, fuel moisture, and atmospheric conditions all play important roles in reducing air quality concerns, ensuring a balanced approach that corresponds with environmental goals while protecting public health (Rafaj et al., 2018). Prescribed burning has effects on carbon sequestration in ecosystems. Controlled burns promote the establishment of fire-adapted plants and avoid uncontrolled wildfires, resulting in a sustainable carbon footprint (Lipsett-Moore et al., 2018; Simões et al., 2023). Prescribed burning is most successful when it is integrated into comprehensive land management schemes (Neidermeier et al., 2023). Controlled burn solutions aim to balance ecological, social, and economic advantages with conservation aims (Pais et al., 2023; Simões et al., 2023).

S.3.3 Options for future changes to fire management

The ABC-iCAP project has produced fire emissions “storylines” for different fire management, natural resource and fire fighting policies, and socioeconomic drivers of fire activity for the countries that comprise the Arctic Council, with a focus on Arctic, boreal, and temperate ecosystems (Paunu et al., 2023). The primary aim is that these future Arctic and boreal fire storylines will be integrated with SSP-aligned emission scenarios and SSP emission estimates and modeling efforts. Three storylines were developed for each of the eight Arctic states, representing low fire activity (dubbed the “We Got This” storyline), high fire activity (“Let It Burn” storyline), and middle-of-the-road fire activity (“The Fire Will Come” storyline). A natural extension of these Fire Management storylines is the product of a spatiotemporally explicit emissions dataset that can be used in future climate

modeling efforts. Further, the process by which the storylines was developed, including country- and ecosystem-specific management and policy variables, could be replicated globally via local experts. The future fire management scenarios and needed emission datasets are unfortunately not available in time for this project, but the specific need to include fire management, fire policies, and socioeconomic influences on global fire activity is a parallel effort required to produce more accurate modeling of global fire emissions.

S4. Inverse modelling of atmospheric transport

Observations of atmospheric concentrations can provide information on sources and sinks in the atmosphere. As mentioned previously, gases concentrations can have their surface fluxes either modified by atmospheric transport or by chemical reactions or physical processes in the atmosphere. By combining a chemistry transport model (CTM) and observed atmospheric gas mole fractions, we can infer surface-atmosphere gas emissions with an atmospheric inversion (also called top-down approach, Bolin and Keeling, 1963). Inversions consist in finding a set of statistically optimal fluxes satisfying measurements and prior inventories (or bottom-up emission inventories) within their respective uncertainties. As an inverse problem, the upwind gas fluxes are estimated from the downwind observed gas mole fractions. The surface gas fluxes are adjusted so that forward-simulated gas mole fractions better match the gas measurements while considering the uncertainty statistics on the observations, transport, and prior surface fluxes. Hence, with inversions, the CTM are better constrained from observed concentration than with forward simulation alone.

Particularly, with dense satellite constellations and networks of surface observations, data assimilation and so inversion is commonly used to quantify emissions. Inverse modeling has been largely used and applied to different scale problems, going from locale and regional emissions (Vautard et al., 2000, Byrne et al., 2022), to optimize global estimations (Houweling et al., 1999, Arellano et al., 2006, Peiro et al., 2022a, 2022b).

Inverse problem are ill-posed meaning that the solution is underdetermined by the observational constraints, which is why additional information are required, such as prior emissions and their uncertainties. This is performed using Bayesian techniques to produce a unique solution. If the amount of observation used in the inversion is high, then the posterior emissions will be more strongly impacted by the assimilated data. However, for regions with low density observations, the posterior fluxes will remain similar to the prior fluxes. This is why in recent inversions, satellite measurements are generally used in addition to ground based observations (Peiro et al., 2022a).

Although CTM performances have improved, there are still uncertainties due to complex interplay of source distributions, transport, and chemistry. Inverse modeling hence allows for a better quantification of the variables driving the physical atmospheric system. If prior error settings are posed appropriately, optimizing emissions and concentrations allows the data assimilation system to address issues on transport or observing errors (Schuh et al., 2019, 2022). As described in the model intercomparison projects (MIPs), the use of multiple inversions performing using different CTM and prior constraints, can help to partially account for systematic errors related to model transport and prior information (Crowell et al., 2019, Peiro et al., 2022a). Even though variability among transport models remains the largest source of uncertainty across global flux inversion

systems, Schuh et al., (2019) suggest the importance of using model ensembles and independent observations for evaluation.

Table S1: Observations available for model evaluation.

Observation	time period	region	species	link to dataset if publicly available	link to information about the dataset	Comments
FIREX-AQ campaign	Summer 2019	Western North America		https://csl.noaa.gov/projects/firex-aq/ ; https://agupubs.onlinelibrary.wiley.com/doi/10.1029/2022JD037758	https://csl.noaa.gov/projects/firex-aq/	aircraft and surface measurements
COALA: Characterizing Organics and Aerosol Loading over Australia	Jan to March 2020	Australia	aerosols, NO _x , O ₃ , CO, VOCs, BVOCs (e.g., isoprene, monoterpenes and sesquiterpenes)		https://www2.acom.ucar.edu/campaigns/coala	COALA white paper: https://www2.acom.ucar.edu/sites/default/files/documents/COALA_whitepaper_v20180410.pdf
TOAR tropospheric ozone database	Long-term, but depends on each site	global	ozone and precursors	https://toar-data.org/	https://toar-data.org/	surface, satellite, aircraft, ozonesondes
The Western Wildfire Experiment for Cloud Chemistry, Aerosol Absorption and Nitrogen (WE-CAN) field campaign	Summer 2018	Western U.S.	CO, CO ₂ , CH ₄ , N ₂ O, H ₂ O, NO, NO ₂ , O ₃		https://www2.acom.ucar.edu/lab-annual-report/2018/1c-western-wildfire-experiment-cloud-chemistry-aerosol-absorption-and-nitrogen-we-can	
DOE Layered Atlantic Smoke Interactions with Clouds (LASIC) campaign	June 2016-October 2017	Ascension Island	SP2, ACSM, optical properties	ARM archive: https://www.arm.gov/research/campaigns/armf2016lasic	Zuidema et al. 2018	surface measurements only
NASA Observations of Aerosols above CLouds and their interactionS	September 2016, August 2017,	Namibia (2016), Sao Tome (2017, 2018)	see Redemann et al. 2021 ACP overview paper	https://doi.org/10.5067/Suborbital/ORACLES		

(ORACLES) aircraft campaign	October 2018		https://doi.org/10.5194/acp-21-1507-2021			
Cloud-Aerosol-Radiation Interaction and Forcing - Year 2017 (CLARIFY-2017) UK aircraft campaign	August-September 2017	Ascension Island (8S, 14.5W)	see Haywood et al. 2021 ACP overview paper	British Atmospheric Data Center (BADC) https://data.ceda.ac.uk/badc/faam/data	Haywood et al. 2021	ORACLES, LASIC, CLARIFY publications collected into an ACP special issue
Organic aerosol observation campaign in East Asia	December 2009–November 2011	Rishiri (45.2N, 141.3E), Okinawa (26.9N, 128.2E), Japan	Carbonaceous (OC, EC), some organic tracers (e.g., levoglucosan, mannitol, SOA tracers), inorganic ions.		https://doi.org/10.5194/acp-15-1959-2015 , https://doi.org/10.1016/j.envpol.2019.01.003	surface measurements for aerosol chemical components
Stockholm Convention Global Monitoring Plan Data Warehouse	Depending on reporting programs (up to 2020)	Global	POPs	https://www.pops-gmp.org/		Annual averages (all national programs contribute to this data warehouse, e.g. GAPS, GLB, EMEP, AMAP etc.)
AGES+ campaign	Summer 2023	North America			https://csl.noaa.gov/projects/ages/	multiple aircraft measurements
MOYA campaign	2017 and 2019	Africa			https://acp.copernicus.org/articles/20/15443/2020/	
BORTAS campaign	2010-2011	North America and Europe	tropospheric oxidants		https://data.ceda.ac.uk/badc/bortas	ACP Special issue: Quantifying the impact of Boreal fires on tropospheric oxidants over the Atlantic using aircraft and satellites (BORTAS) (copernicus.org) Palmer et al. (2013)
WE-CAN campaign	2018					

Table S2. Summary of historical meteorological datasets/reanalyses.

reanalysis	organization	years	Temporal resolution	Vertical resolution	Horizontal resolution	reference
MERRA-2	NASA's Global Modelling and Assimilation Office and distributed by the Goddard Earth Sciences Data and Information Services Center	1980 to the present	hourly	42 pressure vertical levels	0.625° x 0.5° (latitude x longitude)	Gelaro et al., 2017
ERA5	ECMWF	1940 to the present	hourly	37 vertical pressure levels + optional 137 sigma levels	~31 km	Hersbach et al. 2020
NCEP/NCAR Reanalysis 1	NCEP, NCAR	1948 to the present	6-hourly, daily, monthly	17 pressure levels, and 28 sigma levels	2.5° x 2.5°	Kalnay et al., 1996
JRA-55	Japan Meteorological Agency	1958 to the present	3-hourly	60 vertical levels up to 0.1 hPa	TL319	Kobayashi et al., 2015

Table S3. Widely used quantitative Health Risk Assessment tools (Adapted from: Hassan Bhat et al., 2021)

S. No.	Tool	Developer	Reference
1.	Air Quality (Air Q+)	World Health Organization (WHO)	WHO, 2016; Goel et al., 2021
2	BenMAP-CE	United States Environmental Protection Agency (USEPA)	Sacks et al., 2018; 2020
3	The Simple Interactive Model for better Air quality (SIM-air)	Urban Emissions	Guttikunda and Jawahar, 2012; Guttikunda and Calori 2009
4	EcoSense	Institute of Energy Economics and Rational Energy Use (IER), University of Stuttgart	Wagner et al., 2013; Weichenthal et al., 2014

5	Greenhouse gas—Air pollution Interactions and Synergies (GAINS) model	International Institute for Applied Systems Analysis (IIASA)	Nguyen et al., 2008; Liu et al., 2013
6	CO-Benefits Risk Assessment (COBRA) Health Impacts Screening and Mapping Tool	The United States Environmental Protection Agency (EPA)	USEPA, 2021; Mailloux et al., 2022
7	Aphekom	French Institute of Public Health Surveillance	Henschel et al., 2013; Pascal et al., 2013
8	BaPeq and Incremental Lifetime Cancer Risk (ILCR)	The United States Environmental Protection Agency (EPA)	USEPA, 2005; Chen and Liao, 2006

Table S4. Comparative analysis between the different Air pollution Health Risk Assessment tools (Adapted from: Hassan Bhat et al., 2021)

Characteristic	AIRQ2.2	BenMAP-CE	COBRA	ILCR	SIM-Air	GAINS	EcoSense
Health Impacts							
Mortality (cases)	✓	✓	✓	✓	✓	✓	✓
Disability-adjusted life years (DALY)	✓	✓	✓			✓	✓
Morbidity (cases)	✓	✓	✓	✓	✓		✓
Economic Impacts	✓	✓	✓		✓		✓
Pollutants:							
PM _{2.5}	✓	✓		✓	✓	✓	✓
PM ₁₀	✓	✓			✓	✓	✓
Ozone	✓	✓				✓	✓
NO ₂	✓	✓	✓			✓	✓
SO ₂	✓	✓	✓			✓	✓
CO	✓	✓				✓	✓
Other	Black smoke		VOC	VOCs, PAHs		CO ₂ , VOC, CH ₄ , N ₂ O	Hydrocarbons, dioxins and heavy metals
Spatial Resolution							
Regional	✓	✓	✓	✓	✓	✓	✓
National	✓	✓	✓	✓		✓	
City-level	✓	✓	✓	✓	✓	✓	
Household/Indoor	✓		✓	✓		✓	✓

References

- Aisbett, B., Wolkow, A., Sprajcer, M., and Ferguson, S. A.: “Awake, smoky, and hot”: providing an evidence-base for managing the risks associated with occupational stressors encountered by wildland firefighters, *Appl. Ergon.*, 43, 916–925, 2012.
- Alexaki, N., van den Hof, M., and Jol, K.: From Burning to Buying: Creating A Circular Production Chain Out of Left-Over Crop Residue from Indian Farm Land, Netherlands Enterprise Agency, Utrecht, 1–30, <https://www.rvo.nl/sites/default/files/2019/12/MVO-Nederland-rapport-India.pdf> TS120 (last accessed 20 May 2025), 2019.
- Arellano, A. F., Kasibhatla, P. S., Giglio, L., van der Werf, G. R., Randerson, J. T., Collatz, G. J.: Time-dependent inversion estimates of global biomass-burning CO emissions using Measurement of Pollution in the Troposphere (MOPITT), *J. Geophys. Res.*, 111, D09303, <https://doi.org/10.1029/2005JD006613>, 2006.
- Ascoli, D., Plana, E., Oggioni, S. D., Tomao, A., Colonico, M., Corona, P., Giannino, F., Moreno, M., Xanthopoulos, G., Kaoukis, K., Athanasiou, M., Colaço, M. C., Rego, F., Sequeira, A. C., Acácio, V., Serra, M., and Barbati, A.: Fire-smart solutions for sustainable wildfire risk prevention: Bottom-up initiatives meet top-down policies under EU green deal, *Int. J. Disast. Risk Reduct.*, 92, 103715, <https://doi.org/10.1016/j.ijdrr.2023.103715>, 2023.
- Bories, C., Aouba, L., Vedrenne, E., Vilarem, G.: Fired clay bricks using agricultural biomass wastes: study and characterization, *Constr. Build. Mater.*, 91, 158–163, <https://doi.org/10.1016/j.conbuildmat.2015.05.006>, 2015.
- Bran SH, Macatangay R, Surapipith V, Chotamonsak C, Chantara S, Han Z, Li J (2022) Surface PM2.5 mass concentrations during the dry season over northern Thailand: sensitivity to model aerosol chemical schemes and the effects on regional meteorology. *Atmospheric Research*, 277, 106303; doi:10.1016/j.atmosres.2022.106303.
- Byrne, B., Baker, D. F., Basu, S., Bertolacci, M., Bowman, K. W., Carroll, D., Chatterjee, A., Chevallier, F., Ciais, P., Cressie, N., Crisp, D., Crowell, S., Deng, F., Deng, Z., Deutscher, N. M., Dubey, M. K., Feng, S., García, O. E., Griffith, D. W. T., Herkommer, B., Hu, L., Jacobson, A. R., Janardanan, R., Jeong, S., Johnson, M. S., Jones, D. B. A., Kivi, R., Liu, J., Liu, Z., Maksyutov, S., Miller, J. B., Miller, S. M., Morino, I., Notholt, J., Oda, T., O’Dell, C. W., Oh, Y.-S., Ohyama, H., Patra, P. K., Peiro, H., Petri, C., Philip, S., Pollard, D. F., Poulter, B., Remaud, M., Schuh, A., Sha, M. K., Shiomi, K., Strong, K., Sweeney, C., Té, Y., Tian, H., Velasco, V. A., Vrekoussis, M., Warneke, T., Worden, J. R., Wunch, D., Yao, Y., Yun, J., Zammit-Mangion, A., and Zeng, N.: National CO₂ budgets (2015–2020) inferred from atmospheric CO₂ observations in support of the global stocktake, *Earth Syst. Sci. Data*, 15, 963–1004, <https://doi.org/10.5194/essd-15-963-2023>, 2023.
- Crowell, S., Baker, D., Schuh, A., Basu, S., Jacobson, A. R., Chevallier, F., Liu, J., Deng, F., Feng, L., McKain, K., Chatterjee, A., Miller, J. B., Stephens, B. B., Eldering, A., Crisp, D., Schimel, D., Nassar, R., O’Dell, C. W., Oda, T., Sweeney, C., Palmer, P. I., and Jones, D. B. A.: The 2015–2016 carbon cycle as seen from OCO-2 and the global in situ network, *Atmos. Chem. Phys.*, 19, 9797–9831, <https://doi.org/10.5194/acp-19-9797-2019>, 2019.
- Ditomaso, J. M., Brooks, M. L., Allen, E. B., Minnich, R., Rice, P. M., and Kyser, G. B.: Control of Invasive Weeds with Prescribed Burning, *Weed Technol.*, 20, 535–548, <https://doi.org/10.1614/WT-05-086R1.1>, 2006.
- Domingo, J., De Miguel, E., Hurtado, B., Métayer, N., Bamière, L., Pardon, L., Bochu, J., Pointereau, P., and Pellerin, S.: Measures at farm level to reduce greenhouse gas emissions from EU

- agriculture, Notes, Policy Dep. B Struct. Cohes. Policies, [https://www.europarl.europa.eu/RegData/etudes/note/join/2014/513997/IPOL-AGRI_NT\(2014\)513997_EN.pdf](https://www.europarl.europa.eu/RegData/etudes/note/join/2014/513997/IPOL-AGRI_NT(2014)513997_EN.pdf) (last access 20 May 2025), 2014.
- Gelaro, R., W. McCarty, M. J. Suárez, R. Todling, A. Molod, L. Takacs, C. A. Randles, A. Darmenov, M. G. Bosilovich, R. Reichle, et al.: The Modern-Era Retrospective Analysis for Research and Applications, Version 2 (MERRA-2)., *J. Climate*, 30, 14, doi:10.1175/JCLI-D-16-0758.1, 2017.
- Govardhan, G., Ambulkar, R., Kulkarni, S., Vishnoi, A., Yadav, P., Abida, B., Choudhury, B. A., Khare, M., and Ghude, S. D.: Stubble-burning activities in north-western India in 2021: Contribution to air pollution in Delhi, *Heliyon*, 9, e16939, 85 <https://doi.org/10.1016/j.heliyon.2023.e16939>, 2023.
- Gummert, M., Hung, V. V., Pauline, C., and Douthwaite, B.: Sustainable Rice Straw Management, Springer International Publishing, <https://doi.org/10.1007/978-3-030-32373-8>, 2020
- Guttikunda, S. and Calori, G.: Simplified Atmospheric Transport Modelling System (ATMoS-4.0) for the SIM-air tool, SIM-air Working Paper Series 30-2009, SIAM, TS162, 2009.
- Guttikunda, S. K. and Jawahar, P.: Application of SIM-air modelling tools to assess air quality in Indian cities, *Atmos. Environ.*, 62, 551–561, 2012.
- Hadley, M., Vedanthan, R., and Fuster, V.: Air pollution and cardiovascular disease: a window of opportunity, *Nat. Rev. Cardiol.*, 15, 193–194, <https://doi.org/10.1038/nrcardio.2017.207>, 2018.
- Hassan Bhat, T., Jiawen, G., & Farzaneh, H.: Air pollution health risk assessment (AP-HRA), principles and applications. *International journal of environmental research and public health*, 18, 4, 1935, 2021.
- Haywood, J. M., S. Abel, P. Barrett, et al, 2021: Overview: The CLOUD-Aerosol-Radiation Interaction and Forcing: Year-2017 (CLARIFY-2017) measurement campaign, *Atmos. Chem. Phys.*, 21, p. 1049-1084, doi:10.5194/acp-21-1049-2021
- He, T., Lamont, B. B., and Pausas, J. G.: Fire as a key driver of Earth's biodiversity, *Biol. Rev.*, 94, 1983–2010, <https://doi.org/10.1111/brev.12544>, 2019.
- He, G., Liu, T., and Zhou, M.: Straw burning, PM_{2.5}, and death: Evidence from China, *J. Dev. Econom.*, 145, 102468, doi: [10.1016/j.jdeveco.2020.102468](https://doi.org/10.1016/j.jdeveco.2020.102468), 2020.
- Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, D., Simmons, A., Soci, C., Abdalla, S., Abellan, X., Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M., de Chiara, G., Dahlgren, P., Dee, D., Diamantakis, M., Dragani, R., Flemming, J., Forbes, R., Fuentes, M., Geer, A., Haimberger, L., Healy, S., Hogan, R.J., Hólm, E., Janisková, M., Keeley, S., Laloyaux, P., Lopez, P., Lupu, C., Radnoti, G., de Rosnay, P., Rozum, I., Vamborg, F., Villaume, S., Thépaut, J.N.: The ERA5 global reanalysis. *Q. J. R. Meteorol. Soc.* 146, 1999–2049. <https://doi.org/10.1002/QJ.3803>, 2020.
- Houweling, S., Kaminski, T., Dentener, F., Lelieveld, J., Heimann, M.: Inverse modelling of methane sources and sinks using the adjoint of a global transport model, *J. Geophys. Res.*, 104, 26137–26160, 1999.
- Jain, M., Saxena, P., Sharma, S. and Sonwani, S.: Investigation of forest fire activity changes over the central India domain using satellite observations during 2001–2020. *GeoHealth*, 5, 12, p.e2021GH000528, 2021.
- Jiang, L., Zhang, J., Wang, H. H., Zhang, L., and He, K.: The impact of psychological factors on farmers' intentions to reuse agricultural biomass waste for carbon emission abatement, *J. Clean. Prod.*, 189, 797–804, <https://doi.org/10.1016/j.jclepro.2018.04.040>, 2018.

- Kalnay, E., M. Kanamitsu, R. Kistler, W. Collins, D. Deaven, L. Gandin, M. Iredell, S. Saha, G. White, J. Wooll, Y. Zhu, M. Chelliah, W. Ebisuzaki, W. Higgins, J. Janowiak, K. C. Mo, C. Ropelewski, J. Wang, A. Leetmaa, R. Reynolds, Roy Jenne, and Dennis Joseph: The NCEP/NCAR 40-year reanalysis project, *Bull. Amer. Meteor. Soc.*, 77, 437-470, 1996.
- Kobayashi, S., Y. Ota, Y. Harada, A. Ebita, M. Moriya, H. Onoda, K. Onogi, H. Kamahori, C. Kobayashi, H. Endo, K. Miyaoka, and K. Takahashi: The JRA-55 Reanalysis: General Specifications and Basic Characteristics *J. Met. Soc. Jap.*, 93, 1, 5-48, doi:10.2151/jmsj.2015-001), 2015.
- Kumar, R., Barth, M. C., Pfister, G. G., Nair, V. S., Ghude, S. D., and Ojha, N.: What controls the seasonal cycle of black carbon aerosols in India?, *J. Geophys. Res.-Atmos.*, 120, 7788–7812, <https://doi.org/10.1002/2015JD023298>, 2015.
- Lamont, B. B.: Historical links between climate and fire on species dispersion and trait evolution, *Plant Ecol.*, 223, 711–732, <https://doi.org/10.1007/s11258-022-01232-x>, 2022.
- Lelieveld, J., Evans, J. S., Fnais, M., Giannadaki, D., and Pozzer, A.: The contribution of outdoor air pollution sources to premature mortality on a global scale, *Nature*, 525, 367–371, <https://doi.org/10.1038/nature15371>, 2015.
- Lipsett-Moore, G. J., Wolff, N. H., and Game, E. T.: Emissions mitigation opportunities for savanna countries from early dry season fire management, *Nat. Commun.*, 9, 2247, <https://doi.org/10.1038/s41467-018-04687-7>, 2018.
- Liu, F., Klimont, Z., Zhang, Q., Cofala, J., Zhao, L., Huo, H., Nguyen, B., Schopp, W., Sander, R., Zheng, B., Hong, C., He, K., Amann, M., Heyes, C.: Integrating mitigation of air pollutants and greenhouse gases in Chinese cities: development of GAINS-City model for Beijing, *J. Clean. Prod.*, 58, 25–33, 2013.
- Mailloux, N. A., Abel, D.W., Holloway, T., and Patz, J. A.: Nationwide and regional PM_{2.5}-related air quality health benefits from 35 the removal of energy-related emissions in the United States, *GeoHealth*, 6, e2022GH000603, doi: 10.1029/2022GH000603, 2022
- Masters J.: Reviewing the horrid global 2020 wildfire season. *Yale Climate Connections*, 2021.
- McCarty, J. L., Aalto, J., Paunu, V.-V., Arnold, S. R., Eckhardt, S., Klimont, Z., Fain, J. J., Evangeliou, N., Venäläinen, A., Tchepakova, N. M., Parfenova, E. I., Kupiainen, K., Soja, A. J., Huang, L., and Wilson, S.: Reviews and syntheses: Arctic fire regimes and emissions in the 21st century, *Biogeosciences*, 18, 5053–5083, <https://doi.org/10.5194/bg-18-5053-2021>, 2021.
- Mittal, S. K., Singh, N., Agarwal, R., Awasthi, A., and Gupta, P. K.: Ambient air quality during wheat and rice crop stubble burning episodes, *Atmos Environ.*, 43, 238–244, 2009.
- Molina-Terres, D.M., Xanthopoulos G, Diakakis M, Ribeiro L, Caballero D, Delogu GM, et al.: Analysis of forest fire fatalities in southern Europe: Spain, Portugal, Greece and Sardinia (Italy). *Int J Wildland Fire*, 28, 2, 85–98, <https://doi.org/10.1071/WF18004>, 2019.
- Moran J, NaSuwan C, Poocharoen O-O (2019) The haze problem in Northern Thailand and policies to combat it: a review. *Environmental Science and Policy* **97**: 1–15; doi:10.1016/j.envsci.2019.03.016.
- Neidermeier, A. N., Zagaria, C., Pampanoni, V., West, T. A. P., and Verburg, P. H.: Mapping opportunities for the use of land management strategies to address fire risk in Europe, *J. Environ. Manage.*, 346, 118941, <https://doi.org/10.1016/j.jenvman.2023.118941>, 2023.
- Nguyen, T. B., Schoepp, W., and Wagner, F.: GAINS-BI: business intelligent approach for greenhouse gas and air pollution interactions and synergies information system, in: *Proceedings Of The 10th*

- International Conference On Information Integration And Web-Based Applications & Services, November 2008, 332–338, doi: [10.1145/1497308.1497369](https://doi.org/10.1145/1497308.1497369), 2008.
- ODF – Oregon Department of Forestry: ODF Fire History 1911–2020, State Library of Oregon, Salem, OR, USA, <https://www.oregon.gov/odf/fire/documents/odf-century-fire-history-chart.pdf> (last accessed 20 May 2025), 2020.
- Pais, S., Aquilué, N., Honrado, J. P., Fernandes, P. M., and Regos, A.: Optimizing Wildfire Prevention through the Integration of Prescribed Burning into ‘Fire-Smart’ Land-Use Policies, *Fire*, 6, 457, <https://doi.org/10.3390/fire6120457>, 2023.
- Palmer, P. I., Parrington, M., Lee, J. D., Lewis, A. C., Rickard, A. R., Bernath, P. F., Duck, T. J., Waugh, D. L., Tarasick, D. W., Andrews, S., Aruffo, E., Bailey, L. J., Barrett, E., Bauguutte, S. J.-B., Curry, K. R., Di Carlo, P., Chisholm, L., Dan, L., Forster, G., Franklin, J. E., Gibson, M. D., Griffin, D., Helmig, D., Hopkins, J. R., Hopper, J. T., Jenkin, M. E., Kindred, D., Kliever, J., Le Breton, M., Matthiesen, S., Maurice, M., Moller, S., Moore, D. P., Oram, D. E., O'Shea, S. J., Owen, R. C., Pagniello, C. M. L. S., Pawson, S., Percival, C. J., Pierce, J. R., Punjabi, S., Purvis, R. M., Remedios, J. J., Rotermund, K. M., Sakamoto, K. M., da Silva, A. M., Strawbridge, K. B., Strong, K., Taylor, J., Trigwell, R., Tereszchuk, K. A., Walker, K. A., Weaver, D., Whaley, C., and Young, J. C.: Quantifying the impact of BOREal forest fires on Tropospheric oxidants over the Atlantic using Aircraft and Satellites (BORTAS) experiment: design, execution and science overview, *Atmos. Chem. Phys.*, 13, 6239–6261, <https://doi.org/10.5194/acp-13-6239-2013>, 2013.
- Pascal, M., Corso, M., Chanel, O., Declercq, C., Badaloni, C., Cesaroni, G., Henschel, S., Meister, K., Haluza, D., Martin-Olmedo, P., Medina, S., Aphekom Group: Assessing the public health impacts of urban air pollution in 25 European cities: results of the Aphekom project, *Sci. Total Environ.*, 449, 390–400, 2013.
- Paunu, V.-V., McCarty, J. L., Lipsanen, A., and Entsaló, I.: Fire in the Arctic: Current Trends and Future Pathways, Arctic Black Carbon impacting on Climate and Air Pollution (ABC-iCAP) Project Technical Report 1, November 2023, TS213 , vi C 20 pp., 2023.
- Pausas, J. G. and Keeley, J. E.: Wildfires as an ecosystem service, *Front. Ecol. Environ.*, 17, 289–295, <https://doi.org/10.1002/fee.2044>, 2019.
- Peiro, H., Crowell, S., Schuh, A., Baker, D. F., O'Dell, C., Jacobson, A. R., Chevallier, F., Liu, J., Eldering, A., Crisp, D., Deng, F., Weir, B., Basu, S., Johnson, M. S., Philip, S., and Baker, I.: Four years of global carbon cycle observed from the Orbiting Carbon Observatory 2 (OCO-2) version 9 and in situ data and comparison to OCO-2 version 7, *Atmos. Chem. Phys.*, 22, 1097–1130, <https://doi.org/10.5194/acp-22-1097-2022>, 2022a.
- Peiro, H., Crowell, S., and Moore III, B.: Optimizing 4 years of CO₂ biospheric fluxes from OCO-2 and in situ data in TM5: fire emissions from GFED and inferred from MOPITT CO data, *Atmos. Chem. Phys.*, 22, 15817–15849, <https://doi.org/10.5194/acp-22-15817-2022>, 2022b.
- Phoo WW, Manomaiphiboon K, Jaroornrattanapak N, Yodcum J, Sarinnapakorn K, Bonnet S, Aman N, Junpen A, Devkota B, Wang Y, Wilasang C (2024) Fire activity and fire weather in a Lower Mekong subregion: association, regional calibration, weather-adjusted trends, and policy implications. *Natural Hazards*; doi:10.1007/s11069-024-06743-6. (accepted)
- Psaropoulos J.: Greek wildfires devastated land. They also took away livelihoods. Al Jazeera. [cited 2021 December 9], Sept 20, 2021.
- Rafaj, P., Kiesewetter, G., Gül, T., Schöpp, W., Cofala, J., Klimont, Z., Purohit, P., Heyes, C., Amann, M., Borken-Kleefeld, J., and Cozzi, L.: Outlook for clean air in the context of sustainable

- development goals, *Global Environ. Change*, 53, 1–11, <https://doi.org/10.1016/j.gloenvcha.2018.08.008>, 2018.
- Redemann, J., R. Wood, P. Zuidema, S. Doherty, B. Luna, et al, 2021: An overview of the ORACLES (ObseRvations of Aerosols above CLouds and their intEractionS) project: aerosol-cloud-radiation interactions in the Southeast Atlantic basin. *Atmos. Chem. Phys.*, **21**, p. 1507-1563, doi:[10.5194/acp-21-1507-2021](https://doi.org/10.5194/acp-21-1507-2021)
- Reisen, F., Hansen, D., and Meyer, C. P.: Exposure to bushfire smoke during prescribed burns and wildfires: firefighters' exposure risks and options, *Environ. Int.*, 37, 314–321, 2011.
- Sacks, J. D., Lloyd, J. M., Zhu, Y., Anderton, J., Jang, C. J., Hubbell, B., and Fann, N.: The Environmental Benefits Mapping and Analysis Program–Community Edition (BenMAP–CE): A tool to estimate the health and economic benefits of reducing air pollution, *Environ. Model. Softw.*, 104, 118–129, 2018.
- Saggu, G. S., Mittal, S. K., Agarwal, R., and Beig, G.: Epidemiological study on respiratory health of school children of rural sites of Malwa region (India) during post-harvest stubble burning events, *MAPAN*, 33, 281–295, <https://doi.org/10.1007/s12647-018-0259-3>, 2018.
- Sagra, J., Moya, D., Plaza-Álvarez, P. A., Lucas-Borja, M. E., González-Romero, J., De las Heras, J., Alfaro-Sánchez, R., and Ferrandis, P.: Prescribed fire effects on early recruitment of Mediterranean pine species depend on fire exposure and seed provenance, *Forest Ecol. Manage.*, 441, 253–261, <https://doi.org/10.1016/j.foreco.2019.03.057>, 2019.
- Sahu, L. K. and Saxena, P.: High time and mass resolved PTR-TOFMS measurements of VOCs at an urban site of India during winter: Role of anthropogenic, biomass burning, biogenic and photochemical sources, *Atmos. Res.*, 164, 84–94, 2015.
- Saxena, P., Sonwani, S., Srivastava, A., Jain, M., Srivastava, A., Bharti, A., Rangra, D., Mongia, N., Tejan, S., and Bhardwaj, S.: Impact of crop residue burning in Haryana on the air quality of Delhi, India, *Heliyon*, 7, 5, e06973, 2021.
- Schuh, A. E., Jacobson, A. R., Basu, S., Weir, B., Baker, D., Bowman, K., Chevallier, F., Crowell, S., Davis, K. J., Deng, F., Denning, S., Feng, L., Jones, D. B. A., Liu, J., and Palmer, P. I.: Quantifying the impact of atmospheric transport uncertainty on CO₂ surface flux estimates, *Global Biogeochem. Cy.*, 33, 484–500, <https://doi.org/10.1029/2018GB006086>, 2019.
- Schuh, A. E., Byrne, B., Jacobson, A. R., Crowell, S. M. R., Deng, F., Baker, D. F., Johnson, M. S., Philip, S., and Weir, B.: On the role of atmospheric model transport uncertainty in estimating the Chinese land carbon sink, *Nature*, 603, E13–E14, <https://doi.org/10.1038/s41586-021-04258-9>, 2022.
- Simões, R. S., Ribeiro, P. F., and Santos, J. L.: Estimating the Trade-Offs between Wildfires and Carbon Stocks across Landscape Types to Inform Nature-Based Solutions in Mediterranean Regions, *Fire*, 6, 397, <https://doi.org/10.3390/fire6100397>, 2023.
- Smith, P., Martino, D., Cai, Z., Gwary, D., Janzen, H., Kumar, P., McCarl, B., Ogle, S., O'Mara, F., and Rice, C.: Policy and technological constraints to implementation of greenhouse gas mitigation options in agriculture, *Agr. Ecosyst. Environ.*, 118, 6–28, <https://doi.org/10.1016/j.agee.2006.06.006>, 2007.
- Tanpipat V, Honda K, Nuchaiya P (2009) MODIS hotspot validation over Thailand. *Remote Sensing* 1: 1043–1054; doi:[10.3390/rs1041043](https://doi.org/10.3390/rs1041043).
- Tatro, T. and P. Zuidema (submitted, July 2024): More biomass burning aerosol is being advected westward over the southern tropical Atlantic since 2003. *Science Tot. Env.*, STOTEN-D-24-29649, preprint at doi:[10.31223/X5PD95](https://doi.org/10.31223/X5PD95)

- Thacker, F. E. N., Ribau, M. C., Bartholomeus, H., and Stoof, C. R.: What is a fire resilient landscape? Towards an integrated definition, *Ambio*, 52, 1592–1602, <https://doi.org/10.1007/s13280-023-01891-8>, 2023.
- Thao>NNL, Pimonsree S, Prueksakorn K, Thao PBT, Vongruang P (2022) Public health and economic impact assessment of PM2.5 from open biomass burning over countries in mainland Southeast Asia during the smog episode. *Atmospheric Pollution Research* 13: 101418; doi:10.1016/j.apr.2022.101418.
- Unapumnuk and Yensrong: CLEAR Sky Strategy Joint-Plan of Implementation. Conference Proceedings: Environmental Pollutants and Toxicants Affecting Health: Collaborative Efforts for Improving Quality of Life; Session 2: Air Pollution Solutions for the Transboundary Haze Issue in the Mekong Subregion, 2024.
- van der Werf, G. R., Randerson, J. T., Giglio, L., Collatz, G. J., Mu, M., Kasibhatla, P. S., Morton, D. C., DeFries, R. S., Jin, Y., and van Leeuwen, T. T.: Global fire emissions and the contribution of deforestation, savanna, forest, agricultural, and peat fires (1997–2009), *Atmos. Chem. Phys.*, 10, 11707–11735, <https://doi.org/10.5194/acp-10-11707-2010>, 2010.
- Vautard, R., Beekmann, M., Menut, L., and Lattuati, M.: Applications of adjoint modelling in urban air pollution, *Environ. Model. Softw.*, 15, 703–709, [https://doi.org/10.1016/S1364-8152\(00\)00058-X](https://doi.org/10.1016/S1364-8152(00)00058-X), 2000.
- Volkova, L., Roxburgh, S. H., and Weston, C. J.: Effects of prescribed fire frequency on wildfire emissions and carbon sequestration in a fire adapted ecosystem using a comprehensive carbon model, *J. Environ. Manage.*, 290, 112673, <https://doi.org/10.1016/j.jenvman.2021.112673>, 2021.
- Wagner, F., Heyes, C., Klimont, Z., and Schöpp, W.: The GAINS optimization module: Identifying cost-effective measures for improving air quality and short-term climate forcing, International Institute for Applied Systems Analysis (IIASA), <https://www.jstor.org/stable/resrep15788.3> (last accessed 20 May 2025), 2013.
- Weichenthal, S., Villeneuve, P. J., Burnett, R. T., Van Donkelaar, A., Martin, R. V., Jones, R. R., DellaValle, C. T., Sandler, D. P., Ward, M. H., Hoppin, J. A.: Long-term exposure to fine particulate matter: association with nonaccidental and cardiovascular mortality in the agricultural health study cohort, *Environ. Health Perspect.*, 122, 609–615, 2014.
- Yadav, R.S. Stubble Burning: A Problem for the Environment, Agriculture, and Humans. Down to Earth, <https://www.downtoearth.org.in/blog/agriculture/stubble-burning-a-problem-for-the-environment-agriculture-and-humans-64912>, June 4, 2019.
- Zuidema, P., A. Sedlacek, C. Flynn, S. Springston, R. Delgadillo, J. Zhang, A. Aiken and P. Muradyan, 2018: The Ascension Island boundary layer in the remote southeast Atlantic is often smoky. *Geophys. Res. Lett.*, 45, pp. 4456–4465 doi:[10.1002/2017GL076926](https://doi.org/10.1002/2017GL076926)